

## Differential Geometry III – Homework 7

Submission: 13. Dezember 2024, until 8:15 am (start of the exercise class).

---

### 1. Exercise

(4 points)

The four Kepler-Poinsot polyhedra, which can be understood as generalizations of the platonic solids, are depicted in Figure 1.

- i) Which of the fractional Schläfli symbols  $\{5/2, 3\}$ ,  $\{3, 5/2\}$ ,  $\{3/2, 5\}$  and  $\{5, 3/2\}$  correspond to which Kepler-Poinsot polyhedra and how are these fractional Schläfli symbols calculated?
- ii) The Poinsot polyhedra  $\{3, 5/2\}$  and  $\{5, 3/2\}$  each correspond to a branched covering of the sphere. Determine the degree (= number of sheets) of these coverings and calculate the genus of the covering spaces using the Riemann-Hurwitz formula.




Figure 1: From left to right: The great dodecahedron, the small stellated dodecahedron, the great icosahedron and the great stellated dodecahedron. The graphics in this Figure were created using Robert Webb's software Stella from the webpage <https://www.software3d.com/Stella.php>.

*Please turn the page!*

## 2. Exercise

(4 points)

The sphere inversion is the mapping defined by

$$\mathbb{R}^3 \setminus \{0\} \ni (y_1, y_2, y_3) \mapsto \frac{(y_1, y_2, y_3)}{y_1^2 + y_2^2 + y_3^2}.$$

- i) Determine the isometry of  $\mathbb{S}^3 \setminus \{e_4, -e_4\}$  that corresponds to the inversion at the sphere via the stereographic projection  $\widehat{\text{st}}_3: \mathbb{S}^3 \setminus \{e_4, -e_4\} \rightarrow \mathbb{R}^3 \setminus \{0\}$ .
- ii) Sketch a branched 2-sheeted covering from the surface  $M$  in Figure 2 onto the sphere such that it has a branch point with winding number 2 at every vertex of the spherical tiling that corresponds to a cube.
 

*Hint:* Find a set of 8 points that is invariant under the indicated symmetries of  $M$ . Then, using these points as vertices, draw a quad layout<sup>I</sup> on  $M$  and enumerate the vertices/quads corresponding to the vertices/tiles of the cubic spherical tiling.
- iii) Verify by explicit calculation that the Riemann-Hurwitz formula is satisfied for the branched covering from Part ii).




Figure 2: **Left:** A compact, closed Riemann surface  $M$  of genus 3 that has tetrahedral symmetry and is, in addition, invariant under sphere inversion. The vectors  $e_1$ ,  $e_2$  and  $e_3$  are indicated in red, green and blue, respectively. The unit sphere is indicated by some latitudinal and some longitudinal lines. **Right:** The spherical tiling that corresponds to the cube.

Total: 8

<sup>I</sup>A quad means a simply connected domain with four smooth curve segments (possibly geodesics) as boundary.