

### III. Matching


- $G = (V, E)$  undirected graph
- *Matching*: Subset of edges  $M \subseteq E$ , no two of which share an endpoint.
- *Maximum (cardinality) matching*: Matching of maximum cardinality
- *Perfect matching*: Every vertex in  $V$  is matched.
- *Maximum weighted matching*:  
Given a weight function  $w : E \rightarrow \mathbb{R}$ , find a matching  $M$  such that  $w(M) = \sum_{e \in M} w(e)$  is maximal.

#### Augmenting paths

- Let  $M$  be a matching in  $G = (V, E)$ .
- A path  $P = (v_0, v_1, \dots, v_t)$  in  $G$  is called *M-augmenting* if:
  - $t$  is odd,
  - $v_1 v_2, v_3 v_4, v_{t-2} v_{t-1} \in M$ ,
  - $v_0, v_t \notin \bigcup_{e \in M} e$ .
- If  $P$  is an *M-augmenting* path and  $E(P)$  the edge set of  $P$ , then

$$M' = M \triangle E(P) = (M \setminus E(P)) \cup (E(P) \setminus M)$$

is a matching in  $G$  of size  $|M'| = |M| + 1$ .



#### Berge's Theorem

**Theorem** (Berge 1957)

Let  $M$  be a matching in the graph  $G = (V, E)$ . Then either  $M$  is a maximum cardinality matching or there exists an *M-augmenting* path.

*Generic Matching Algorithm*

*Initialization*:  $M \leftarrow \emptyset$

*Iteration*: If there exists an *M-augmenting* path  $P$ , replace  $M \leftarrow M \triangle E(P)$ .

~> how can one find an *M-augmenting* path?

- Difficult in general ~> Edmonds' matching algorithm (Edmonds 1965)
- Easy for bipartite graphs

#### Bipartite graphs

A graph  $G = (V, E)$  is *bipartite* if there exist  $A, B \subseteq V$  with  $A \cup B = V$ ,  $A \cap B = \emptyset$  and each edge in  $E$  has one end in  $A$  and one end in  $B$ .

**Proposition**

A graph  $G = (V, E)$  is bipartite if and only if each circuit of  $G$  has even length.

## Bipartite matching

Matching augmenting algorithm for bipartite graphs

*Input:* Bipartite graph  $G = (A \cup B, E)$  with matching  $M$ .

*Output:* Matching  $M'$  with  $|M'| > |M|$  or proof that no such matching exists.

*Description:* Construct a directed graph  $D_M$  with the same node set as  $G$ .

For each edge  $e = \{a, b\}$  in  $G$  with  $a \in A, b \in B$ :

if  $e \in M$ , there is the arc  $(b, a)$  in  $D_M$ .

if  $e \notin M$ , there is the arc  $(a, b)$  in  $D_M$ .

Let  $A_M = A \setminus \cup M$  and  $B_M = B \setminus \cup M$ .


$M$ -augmenting paths in  $G$  correspond to directed paths in  $D_M$  starting in  $A_M$  and ending in  $B_M$ .

### Theorem

A maximum-cardinality matching in a bipartite graph  $G = (V, E)$  can be found in time  $O(|V||E|)$ .

## Bipartite matching as a maximum flow problem

- Add a source  $s$  and edges  $(s, a)$  for  $a \in A$ , with capacity 1.
- Add a sink  $t$  and edges  $(b, t)$  for  $b \in B$ , with capacity 1.
- Direct edges in  $G$  from  $A$  to  $B$ , with capacity 1.




- Integral flows  $f$  correspond to matchings  $M$ , with  $\text{val}(f) = |M|$ .
- Ford-Fulkerson takes time  $O(nm)$ , since  $v^* \leq n$ .
- Can be improved to  $O(\sqrt{n}m)$  (Hopcroft-Karp 1973).

## Marriage theorem


**Theorem** (Hall 1935)

A bipartite graph  $G = (A \cup B, E)$ , with  $|A| = |B| = n$ , has a perfect matching if and only if for all  $B' \subseteq B$ ,  $|B'| \leq |N(B')|$ , where  $N(B')$  is the set of all neighbors of nodes in  $B'$ .




**Proof**

- Let  $(S, T)$  be an  $(s, t)$ -cut in the corresponding network.
- Define  $A_S = A \cap S, A_T = A \cap T, B_S = B \cap S, B_T = B \cap T$ .
- Show  $\text{cap}(S, T) \geq n$  (Exercise)
- By the max-flow min-cut theorem, the maximum flow is at least  $n$ .

**Network connectivity: Menger's theorems**

- $G = (V, E)$  directed graph,  $s, t \in V, s \neq t$  non-adjacent.
- **Theorem** (Menger 1927) The maximum number of *arc-disjoint* paths from  $s$  to  $t$  equals the minimum number of arcs whose removal disconnects all paths from  $s$  to  $t$ .
- **Theorem** (Menger 1927) The maximum number of *node-disjoint* paths from  $s$  to  $t$  equals the minimum number of nodes (different from  $s$  and  $t$ ) whose removal disconnects all paths from  $s$  to  $t$ .

**References and further reading**

- A. Schrijver: A Course in Combinatorial Optimization, CWI Amsterdam, 2010, <http://homepages.cwi.nl/~lex/files/dict.pdf>
- K. Mehlhorn: Data Structures and Efficient Algorithms, Vol. 2: Graph Algorithms and NP-Completeness, Springer, 1986, <http://www.mpi-sb.mpg.de/~mehlhorn/DatAlgbooks.html>
- S. Krumke and H. Noltemeier: Graphentheoretische Konzepte und Algorithmen. Teubner, 2005
- R. K. Ahuja, T. L. Magnanti and J. L. Orlin: Network flows. Prentice Hall, 1993