

Prof. Dr. Knut Reinert,
Prof. Dr. Alexander Bockmayr,
René Rahn,
Annika Röhl

November 14, 2014

Algorithms

WS 2014/15

Exercises 5

1. "sparse" skip list (Niveau I)

Each node in the skip list has up to two incoming directed edges pointing to other nodes in the skip list.

- (a) Which edges are really necessary for a search and which can be removed?
- (b) Can you give a rough estimate for the expected number of edges that can be removed?

2. Skip lists (Niveau II)

Proof that the height of a skip list has expected value $O(\log n)$ with high probability, i.e. show that the probability that the height deviates from $\log n$ by a large factor is very low.

Hint: You do not need Chernoff bounds or Markov's inequality to show this.

3. Expected values (Niveau I)

Let X and Y be random variables:

- (a) Prove that $E(X + Y) = E(X) + E(Y)$.
- (b) Assume that X and Y are independent. Prove that $E(XY) = E(X)E(Y)$
- (c) Assume that X takes values $\{0, 1, 2, \dots\}$. Show that $E(X) = \sum_{k=1}^{\infty} \Pr(X \geq k)$.

4. Tree decomposition (Niveau I)

Prove the following theorem:

Let $G = (V, E)$ be a graph, T be a tree decomposition of G , and (x, y) an edge in T . The deletion of (x, y) divides T into two components X and Y . Let V_x and V_y be the 'pieces' of x and y , respectively. Then deleting the set $V_x \cap V_y$ from V disconnects G into the two subgraphs $G_X - (V_x \cap V_y)$ and $G_Y - (V_x \cap V_y)$.

(G_M for $M = X, Y$ is the subgraph of G that consists of all nodes in the 'pieces' of M .)