

# Graph Algorithms

## I. Shortest paths

- $D = (V, A)$  directed graph,  $s, t \in V$ .
- A *walk* is a sequence  $P = (v_0, a_1, v_1, \dots, a_k, v_k)$ ,  $k \geq 0$ , where  $a_i$  is an arc from  $v_{i-1}$  to  $v_i$ , for  $i = 1, \dots, k$ .
- $P$  is a *path*, if  $v_0, \dots, v_k$  are all different.
- If  $s = v_0$  and  $t = v_k$ ,  $P$  is a *s-t walk* resp. *s-t path of length k* (i.e., each arc has length 1).
- The *distance* from  $s$  to  $t$  is the minimum length of any *s-t path* (and  $+\infty$  if no *s-t path* exists).

### Shortest paths with unit lengths

**Algorithm** (Breadth-first search)

*Initialization:*  $V_0 = \{s\}$

*Iteration:*  $V_{i+1} = \{v \in V \setminus (V_0 \cup V_1 \cup \dots \cup V_i) \mid (u, v) \in A, \text{ for some } u \in V_i\}$ ,  
until  $V_{i+1} = \emptyset$ .

*Running time:*  $O(|A|)$

- $V_i$  is the set of nodes with distance  $i$  from  $s$ .
- The algorithm computes shortest paths from  $s$  to all reachable nodes.
- Can be described by a directed tree  $T = (V', A')$  with root  $s$  such that each  $u-v$  path in  $T$  is a shortest  $s-t$  path in  $D$ .

### Shortest paths with non-negative lengths

- Length function  $l : A \rightarrow \mathbb{Q}_+ = \{x \in \mathbb{Q} \mid x \geq 0\}$
- For a walk  $P = (v_0, a_1, v_1, \dots, a_k, v_k)$  define  $l(P) = \sum_{i=1}^k l(a_i)$ .

**Algorithm** (Dijkstra 1959)

*Initialization:*  $U = V$ ,  $f(s) = 0$ ,  $f(v) = \infty$ , for  $v \in V \setminus \{s\}$

*Iteration:* Find  $u \in U$  with  $f(u) = \min\{f(v) \mid v \in U\}$ .


For all  $a = (u, v) \in A$  with  $f(v) > f(u) + l(a)$  let  $f(v) = f(u) + l(a)$ .

Let  $U \leftarrow U \setminus \{u\}$ , until  $U = \emptyset$ .

Upon termination,  $f(v)$  gives the length of a shortest path from  $s$  to  $v$ .

*Running time:*  $O(|V|^2)$  (can be improved to  $O(|A| + |V| \log |V|)$ .)

### Example



Iteration	$u$	$U$	$f[1]$	$f[2]$	$f[3]$	$f[4]$	$f[5]$
0	–	{1, 2, 4, 3, 5}	0	$\infty$	$\infty$	$\infty$	$\infty$
1	1	{2, 3, 4, 5}	0	1	$\infty$	3	10
2	2	{3, 4, 5}	0	1	6	3	10
3	4	{3, 5}	0	1	5	3	9
4	3	{5}	0	1	5	3	6
5	5	{}	0	1	5	3	6

### Application: Longest common subsequence

- Sequences  $a = a_1, \dots, a_m$  and  $b = b_1, \dots, b_n$
- Find the longest common subsequence of  $a$  and  $b$  (obtained by removing symbols in  $a$  or  $b$ ).

*Modeling as a shortest path problem*

- Grid graph with nodes  $(i, j), 0 \leq i \leq m, 0 \leq j \leq n$ .
- Horizontal and vertical arcs of length 1.
- Diagonal arcs  $((i-1, j-1), (i, j))$  of length 0, if  $a_i = b_j$ .

The diagonal arcs on a shortest path from  $(0, 0)$  to  $(m, n)$  define a longest common subsequence.

### Circuits of negative length

- Consider arbitrary length functions  $l : A \rightarrow \mathbb{Q}$ .
- A *directed circuit* is a walk  $P = (v_0, a_1, v_1, \dots, a_k, v_k)$  with  $k \geq 1$  and  $v_0 = v_k$  such that  $v_1, \dots, v_k$  and  $a_1, \dots, a_k$  are all different.
- If  $D = (V, A)$  contains a directed circuit of negative length, there exist  $s$ - $t$  walks of arbitrary small negative length.

#### Proposition

Let  $D = (V, A)$  be a directed graph without circuits of negative length.

For any  $s, t \in V$  for which there exists at least one  $s$ - $t$  walk, there exists a shortest  $s$ - $t$  walk, which is a path.

### Shortest paths with arbitrary lengths

$D = (V, A), n = |V|, l : A \rightarrow \mathbb{Q}$ .

**Algorithm** (Bellman-Ford 1956/58)

Compute  $f_0, \dots, f_n : V \rightarrow \mathbb{R} \cup \{\infty\}$  in the following way:


*Initialization:*  $f_0(s) = 0, f_0(v) = \infty$ , for  $v \in V \setminus \{s\}$

*Iteration:* For  $k = 1, \dots, n$  and all  $v \in V$ :

$$f_k(v) = \min\{f_{k-1}(v), \min_{(u,v) \in A} (f_{k-1}(u) + l(u, v))\}$$

*Running time:*  $O(|V||A|)$

### Example



Iteration $k$	$f_k[1]$	$f_k[2]$	$f_k[3]$	$f_k[4]$	$f_k[5]$
0	0	$\infty$	$\infty$	$\infty$	$\infty$
1	0	7	$\infty$	$\infty$	6
2	0	7	2	4	6
3	0	7	2	4	2
4	0	7	-2	4	2

### Properties

- For each  $k = 0, \dots, n$  and each  $v \in V$ :

$$f_k(v) = \min\{l(P) \mid P \text{ is an } s\text{-}v \text{ walk traversing at most } k \text{ arcs}\}$$

(by induction)

- If  $D$  contains no circuits of negative length,  $f_{n-1}(v)$  is the length of a shortest path from  $s$  to  $v$ .

### Finding an explicit shortest path

- When computing  $f_0, \dots, f_n$  determine a predecessor function  $p : V \rightarrow V$  by setting  $p(v) = u$  whenever  $f_{k+1}(v) = f_k(u) + l(u, v)$ .
- At termination,  $v, p(v), p(p(v)), \dots, s$  gives the reverse of a shortest  $s$ - $v$  path.

#### Theorem

Given  $D = (V, A), s, t \in V$  and  $l : A \rightarrow \mathbb{Q}$  such that  $D$  contains no circuit of negative length, a shortest  $s$ - $t$  path can be found in time  $O(|V||A|)$ .

#### Remark

$D$  contains a circuit of negative length reachable from  $s$  if and only if  $f_n(v) \neq f_{n-1}(v)$ , for some  $v \in V$ .

### NP-completeness

For directed graphs containing circuits of negative length, the problem becomes NP-complete:

#### Theorem

The decision problem

*Input:* Directed graph  $D = (V, A), s, t \in V, l : A \rightarrow \mathbb{Z}, L \in \mathbb{Z}$

*Question:* Does there exist an  $s$ - $t$  path  $P$  with  $l(P) \leq L$ ?

is NP-complete.

#### Corollary

The shortest path problem with arbitrary lengths is NP-complete.

The longest path problem with non-negative lengths is NP-complete.

## Application: Knapsack problem

- Knapsack, volume 8, 5 articles

Article $i$	Volume $a_i$	Value $c_i$
1	5	4
2	3	7
3	2	3
4	2	5
5	1	4

- Objective: Select articles fitting into the knapsack and maximizing the total value.

### Possible models

- *Shortest path model*

- Directed graph with nodes  $(i, x), 0 \leq i \leq 6, 0 \leq x \leq 8$ .
- Arcs from  $(i-1, x)$  to  $(i, x)$  resp.  $(i, x+a_i)$  of length 0 resp.  $-c_i$ , for  $0 \leq i \leq 5$ .
- Arcs from  $(5, x)$  to  $(6, 8)$  of length 0, for  $0 \leq x \leq 6$ .
- A shortest path from  $(0, 0)$  to  $(6, 8)$  gives an optimal solution.

$\rightsquigarrow$  *pseudo-polynomial algorithm*

- *Linear 0-1 model*

$$\max\{4x_1 + 7x_2 + 3x_3 + 5x_4 + 4x_5 \mid 5x_1 + 3x_2 + 2x_3 + 2x_4 + x_5 \leq 8, x_1, \dots, x_5 \in \{0, 1\}\}$$