

Prof. Dr. Alexander Bockmayr,  
Prof. Dr. Knut Reinert,  
Sandro Andreotti

January 15, 2011

## Discrete Mathematics for Bioinformatics (P1)

WS 2010/11

Exercises 10

### 1. Lagrangean Relaxation I (NIVEAU I)

Consider the following problem

$$\begin{array}{rllll} \min & 2x_1 & - & 3x_2 & \\ \text{w.r.t.} & 3x_1 & - & 4x_2 & \geq -6 \\ & -x_1 & + & x_2 & \leq 2 \\ & 6x_1 & + & 2x_2 & \geq 3 \\ & 6x_1 & + & x_2 & \leq 15 \\ & & & x_1, x_2 & \geq 0 \\ & & & x_1, x_2 & \in \mathbb{Z} \end{array}$$

- Draw the corresponding polytope and determine graphically the optimal solution  $Z_{IP}$  of the original problem and  $Z_{LP}$ , the solution of the LP-relaxation.
- Now apply lagrangean relaxation by relaxing the first inequality. Draw the polytope of the relaxed ILP. Determine the set  $X$  of feasible solutions for the relaxed problem.
- The new objective function is then:

$$Z(P) = \min_{(x_1, x_2) \in X} 2x_1 - 3x_2 + p(-6 - 3x_1 + 4x_2)$$

Calculate  $Z_D = \max_{p \geq 0} Z(p)$  and compare this value to  $Z_{IP}$  and  $Z_{LP}$ . (To obtain  $Z_D$ , draw the graphs of the function  $f(p) = 2x_1 - 3x_2 + p(-6 - 3x_1 + 4x_2)$  for all  $(x_1, x_2) \in X$ .)

- repeat a-c for the objective functions  $-x_1 + x_2$  and  $-x_1 - x_2$  and compare  $Z_{LP}$ ,  $Z_D$ , and  $Z_{IP}$ .

## 2. Lagrangean Relaxation II (NIVEAU I)

Prove Lemma 1 (see script page 4001) stating that (in case of a minimization problem) if  $\lambda \geq 0$ , then  $Z(\lambda) \leq Z_{IP}$ , where  $Z_{IP}$  is the optimal value of an original ILP and  $Z(\lambda)$  is the optimal value of the relaxed problem for a given value of the Lagrangean multiplier  $\lambda$ .

## 3. Structural alignment (NIVEAU I)

Prove the two Lemma from the lecture:

**Lemma 1.** *Let  $G = (V, E, H, I)$  be a SEAG with  $n$  alignment edges and  $m$  interaction matches. Then*

- $P_{\mathcal{R}}(G)$  is full-dimensional and
- the inequality  $x_i \leq 1$  is facet-defining iff there is no  $e_j \in E$  in conflict with  $e_i$ .

**Lemma 2.** *Let  $G = (V, E, H, I)$  be a SEAG with  $n$  alignment edges and  $m$  interaction matches. Then*

- The inequality  $x_i \geq 0$  is facet-defining iff  $e_i$  is not contained in an interaction match.
- For each interaction match  $m_{i,j}$  the inequality  $x_{i,j} \geq 0$  is facet-defining.