Duality (2)

Theorem. (Duality Theorem)

If the primal has an optimal solution x^* then the dual has an optimal solution y^* such that

$$c^T x^* = v^{*T} b$$

Proof. (Constructive! → blackboard)

Some properties:

- The dual of the dual is the primal (exercise).
- Corollary: Primal has an optimal solution iff dual has an optimal solution.
- ullet Corollary: Primal unbounded o dual infeasible.
- $\bullet \ \ \text{Corollary: Dual unbounded} \to \text{primal infeasible}.$
- Primal and dual can both be infeasible (Example).

Duality (3)

Practical implications:

- It might be faster to run the Simplex algorithm on the dual.
- Duality provides an elegant way of *proving optimality*. So, solutions come with a certificate, which is always good.

Simplex Algorithm: Geometric view

- Hyperplane $H = \{x \in \mathbb{R}^n \mid a^T x = \beta\}, a \in \mathbb{R}^n \setminus \{0\}, \beta \in \mathbb{R}$
- Closed halfspace $\overline{H} = \{x \in \mathbb{R}^n \mid a^T x \leq \beta\}$
- Polyhedron $P = \{x \in \mathbb{R}^n \mid Ax \leq b\}, A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m$
- Polytope $P = \{x \in \mathbb{R}^n \mid Ax \le b, \ l \le x \le u\}$

The feasible set

$$P = \{x \in \mathbb{R}^n \mid Ax \le b\}$$


of a linear optimization problem is a polyhedron.

Simplex Algorithm: Geometric view (2)

- $P \subseteq \overline{H}, H \cap P \neq \emptyset$ (Supporting hyperplane)
- $F = P \cap H$ (Face)
- dim(F) = 0 (Vertex)
- dim(F) = 1 (Edge)

• $\dim(F) = \dim(P) - 1$ (Facet)

Simplex Algorithm: Geometric view (3)



Simplex Algorithm: Geometric view (4)

Linear optimization problem

$$\max\{c^T x \mid Ax \le b, x \in \mathbb{R}^n, x \ge 0\}$$
 (LP)

Simplex-Algorithm (Dantzig 1947)

- 1. Find a vertex of P.
- 2. Proceed from vertex to vertex along edges of P such that the objective function $z = c^T x$ increases.
- 3. Either a vertex will be reached that is optimal, or an edge will be chosen which goes off to infinity and along which *z* is unbounded.